Use **NCL wipes™ Disinfectant Wipes** in Schools, Colleges, Hospitals, Medical Offices, Nursing homes, Day care centers, Nurseries, Institutional facilities, USDA inspected food processing facilities, Office buildings, Restaurants, Grocery stores, Convenience stores, Hotels, Motels, Prisons, Salons

NCL wipes™ Disinfect Wipes are effective against:

Bacteria

Acinetobacter baumannii
Burkholderia cepacia
Campylobacter jejuni (Campylobacter)
Corynebacterium ammoniagenes
ESBL Escherichia coli
Escherichia coli (E. coli)
Escherichia coli 0157:H7
Klebsiella pneumoniae (NDM-1)
Klebsiella pneumoniae (Klebsiella)
Legionella pneumophilia
Listeria monocytogenes
Salmonella enterica (Salmonella)
Salmonella schottmuelleri (Salmonella)
Salmonella typhi (Salmonella)
Serratia marcescens (Serratia)
Shigella dysenteriae (Shigella)
Pseudomonas aeruginosa (Pseudomonas)
Enterobacter aerogenes
Enterococcus faecalis (Enterococcus)
Enterococcus faecalis - Vancomycin Resistant (VRE)
Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA) (NRS123) (USA400)
Staphylococcus aureus - Community Associated Methicillin-Resistant (CA-MRSA) (NRS384) (USA300)
Staphylococcus aureus - Methicillin-Resistant (MRSA)
Staphylococcus aureus - Multi-Drug Resistant
Staphylococcus aureus (Staph)
Staphylococcus aureus-Vancomycin Intermediate Resistant - (VISA)
Streptococcus pyogenes (Strep)
Vibrio cholerae

Viruses

*2013 Influenza A Virus (H7N9)
*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)
*Herpes Simplex Virus Type 1
*Herpes Simplex Virus Type 2
*HIV-1 (AIDS Virus)
*Human Coronavirus
*Influenza A Virus H1N1
*Influenza Type A Virus / Brazil (the Flu Virus)
*Norwalk Virus (Norovirus)
*Respiratory Syncytial Virus (RSV)
*Rotavirus
*SARS Associated Coronavirus (SARS)
*Vaccinia Virus

Animal Viruses

Avian Influenza Virus H3N2
Avian Influenza Virus H5N1
Canine Distemper Virus
Feline Calicivirus
Newcastle's Disease Virus
Pseudorabies Virus

Fungi

Candida albicans

World Class Cleaning Solutions®

Distributed by:

National Chemical Laboratories, Inc.

401 N. 10th Street • Philadelphia, PA 19123 • (800) 628-2436 • Manufactured in the USA

NCL wipes™

DISINFECTANT WIPES

125 count

Waterfall Fresh

KILLS 99.9% of BACTERIA in 15 SECONDS!

One Step Cleaner / Disinfectant / Deodorizer / Virucide* / Sanitizer

- ***Kills Cold and Flu Virus.*,[1]***
- ***Kills Mold & Mildew.***
- ***Effective in the presence of 5% serum contamination.***
- ***Leaves surfaces* clean and disinfected.***
- ***Does not contain bleach or ammonia.***

A **"One-Step"** disinfectant that is effective against a broad spectrum of bacteria when used as directed.

Kills Pseudomonas, Staph, Salmonella, MRSA, CA-MRSA, E.coli, VRE, Norovirus, HBV, HCV

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride	0.0909%
Dioctyl dimethyl ammonium chloride	0.0364%
Didecyl dimethyl ammonium chloride	0.0545%
Alkyl (C_{14}, 50%; C_{12} 40%; C_{16}, 10%) dimethyl benzyl ammonium chloride	0.1212%
OTHER INGREDIENTS:	**99.6970%**
TOTAL	**100.0000%[1]**

Product No.: 4541

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements and First Aid

NET CONTENTS: 125 WIPES

7" x 8" (17.9 cm x 20.3 cm)

MINIMUM WEIGHT *1.93 lbs (876 g)*

[1] Includes weight of absorbent wipe.

*,[1]On hard, non-porous, non-food surfaces. EPA REG. NO. 6836-340-2296 EPA EST. NO. 70930-PA-001

Use **NCL wipes™ Disinfectant Wipes** on washable hard, nonporous surfaces of: Chairs, Counters, Desks, Doorknobs, Faucets, Glass, Light switches, Showers, Sinks, Tables, Telephones, Toilets, Diaper changing tables, and other hard nonporous surfaces made of Acrylic, Metal, Plastic, Stainless steel, Upholstery-vinyl and Plastic.

TO OPEN PACKAGE:
Open lid. To start wipes, pull up on wipe and move wipe into the small dispensing opening and pull sheet up to remove wipe. The next wipe will pop up automatically. Close lid securely between uses.

DIRECTIONS FOR USE:
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not use as a diaper wipe or for personal cleansing. This is not a baby wipe! Do not flush in toilet.

TO CLEAN AND DEODORIZE: Wipe surface with wipe. Let air dry. Discard used wipe in trash. Do not flush in toilet.

TO CLEAN AND DISINFECT: Wipe surface with wipe until surface is visibly wet. Use enough wipes to thoroughly wet surfaces. Use enough wipes to keep surfaces visibly wet for 4 minutes. Allow surface to remain wet for 4 minutes. For **Candida albicans**, treated surfaces must remain wet for 8 minutes. For **Norovirus and Rotavirus**, treated surfaces must remain wet for 10 minutes. If surfaces are extremely dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush in toilet. Rinse food contact surfaces with clean, potable water after disinfecting. Do not use to disinfect dishes, glassware or utensils.

FOR MOLD AND MILDEW
NCL wipes™ Disinfectant Wipes will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

SANITIZING
To sanitize hard, non-porous, non-food contact surfaces; wipe surface, using enough wipes to allow treated surface to remain wet for 15 seconds for Staphylococcus aureus, Klebsiella pneumoniae, and Campylobacter jejuni. Let surface dry. For highly soiled surfaces clean excess dirt first.

This product is not for use on medical device surfaces.

***KILLS HIV, AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/ BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 or HBV or HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Cleanup must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV, HBV or HCV must be thoroughly cleaned from surfaces and objects before application of **NCL wipes™ Disinfectant Wipes**. **NCL wipes™ Disinfectant Wipes** can be used for this purpose.

Contact time: Leave surface wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses, bacteria and fungi claimed.

Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present , after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children. Keep securely closed. Nonrefillable container. Do not reuse or refill this container. Discard empty container in trash or recycle.

LNW-0001-4541-01

7 52610 94541 6