FOR SAMPLE USE ONLY

READIQUAT

READY-TO-USE DISINFECTANT, CLEANER, VIRUCIDE*.

EPA Reg. No. 10324-85-XXXXX • EPA Est. No. 3838-WI-1

ACTIVE INGREDIENTS:

Alkyl (C_{14} 50%; C_{12} 40%; C_{16} 10%) dimethyl benzyl ammonium chloride	0.034%
Octyl Decyl Dimethyl Ammonium Chloride	0.026%
Didecyl Dimethyl Ammonium Chloride	0.013%
Dioctyl DImethyl Ammonium Chloride	0.013%
INERT INGREDIENTS:	99.914%
TOTAL:	100.000%

CUIDADO. SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

CAUTION. IF YOU CANNOT READ ENGLISH, ASK YOUR SUPERVISOR TO EXPLAIN THE APPROPRIATE DIRECTIONS FOR USE BEFOREWORKING WITH THIS PRODUCT.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel For Additional Precautionary Statements and First Aid Statements.

This formulation is for use on hard, non-porous surfaces in: Institutions, Commercial and Industrial Institutions.

Net Contents: 1 Quart (0.95 Liters)

DISINFECTION PERFORMANCE (in Institutions, Hospitals, Dental Offices, Nursing Homes, and Other Health Care Institutions and facilities, Schools, Restaurants, Food Services, Beverage and Food Processing Plants and Other Non-Medical Facilities): This product kills the following bacteria in 10 minutes in the presence of 5% soil on hard, non-porous surfaces: Burkholderia cepacia, Campylobacter jejuni, Corynebacterium ammoniagenes, Enterobacter aerogenes, Enterobacteriaciae with extended beta-lactamase resistance, Enterobacter cloacae, Enterococcus faecalis, Enterococcus faecium Vancomycin resistant, Escherichia coli, Escherichia coli Antibiotic resistant, Escherichia coli 0157:H7, Klebsiella pneumoniae, Klebsiella pneumoniae Antibiotic resistant, Legionella pneumophila, Listeria monocytogenes, Proteus mirabilis, Proteus vulgaris, Pseudomonas aeruginosa, Pseudomonas aeruginosa Antibiotic resistant, Salmonella enterica, Salmonella schottmuelleri, Salmonella typhi, Serratia marcescens, Shigella dysenteriae, Shigella flexneri, Shigella sonnei, Staphylococcus aureus, Staphylococcus aureus (CA MRSA) Community Acquired Methicillin Resistant, Staphylococcus aureus Methicillin resistant (MRSA), Staphylococcus epidermidis Antibiotic resistant, Streptococcus pyogenes, Vibrio cholerae.

VIRUCIDAL* PERFORMANCE: This product kills the following viruses in 10 minutes in the presence of 5% soil on hard, non-porous surfaces: Avian Influenza A Virus (H5N1), Herpes Simplex Type 1 Virus, Herpes Simplex Type 2 Virus, HIV-1**, Hepatitis C Virus, Hepatitis B Virus, Human Coronavirus, Influenza A/Brazil Virus, Influenza A (H1N1) Virus, Respiratory Syncytial Virus, Vaccinia Virus. ** Indicates a 2-minute contact time is required for this claim.

MILDEWSTATIC PERFORMANCE: This product inhibits the following fungi in 10 minutes in the presence of 5% soil on hard, non-porous surfaces: Aspergillus niger.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISINFECTION/VIRUCIDAL* DIRECTIONS: Pre-clean heavily soiled areas. Apply this undiluted product undiluted to walls, floors and other hard, non-porous surfaces such as Crypton barrier fabric, tables, chairs, countertops, sinks, glazed tile, and glazed porcelain with a brush, cloth, sponge or mechanical spray device so as to thoroughly wet surfaces. If mechanical spray device is used, spray must be coarse. Spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 10 minutes and then let air dry or wipe up excess liquid. Rinse food contact surfaces with potable water after use. Rinsing is not necessary unless floors are to be coated with finish or restorer. Replace solution if it becomes visibly dirty.

This product is not for use on Medical device surfaces.

FOR DISINFECTING HARD, NON-POROUS SURFACES AGAINST AVIAN INFLUENZA A (H5N1) VIRUS: For use in hospitals, hotels, motels and schools, follow the disinfection directions.

***KILLS HIV, HCV AND HBV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1** (HIV-1) associated with AIDS, Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV AND HBV ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Clean-up must always be done wearing protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV-1, HBV & HCV must be thoroughly cleaned from surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** For HIV, let surface remain wet for 2 minutes with undiluted solution. For HBV, HCV and all other organisms listed on label, a 10 minute contact time is required.

FOR USE AS A CLEANER/DISINFECTANT IN FOOD PROCESSING PLANTS: 1. Before using this product in food processing areas, food products and packaging materials must be removed from the room or carefully protected. 2. Apply undiluted solution evenly over surface using a brush, cloth, sponge or mechanical spray device. For sprayer applications, use a coarse spray device. Be sure to wet all surfaces thoroughly. Do not breathe spray. 3. Allow product to remain on surface for ten minutes. 4. Wipe with clean cloth, sponge or paper towel. 5. For heavily soiled areas, thoroughly clean surface prior to disinfecting. 6. When disinfecting food contact surfaces used for food preparation, rinse surfaces thoroughly with potable water. This product must not be used to clean the following surfaces: utensils, glassware and dishes or interior surfaces of processing equipment. 7. Replace solution if it becomes visibly dirty.

TO DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES For countertops, exterior surfaces of appliances, and tables: Before using this product, food products and packaging materials must be removed from area or carefully protected. Pre-clean heavily soiled areas. Apply undiluted solution with a brush, cloth, sponge or hand pump trigger spray device, mechanical spray device so as to wet all surfaces thoroughly. For sprayer applications, use a coarse spray device. Allow surface to remain wet for 10 minutes. Then remove excess liquid and rinse the surface with potable water. Replace solution if it becomes visibly dirty. This product is not for use on dishes, glassware, interior surfaces of appliances or eating utensils or interior surfaces of processing equipment.

DISINFECTION AND CLEANING TOILET BOWLS AND URINALS: 1. Preliminary cleaning is required for heavily soiled surfaces. 2. With a swab applicator, remove water from bowl by forcing out trap. 3. Press swab against side of bowl to remove excess water from applicator. 4. Apply 1-2 oz. of this undiluted product evenly onto surface to be cleaned and disinfected. 5. Swab entire surface area especially under the rim where outlets are located. 6. Allow this product to remain wet on surfaces at least 10 minutes. 7. Flush toilet or urinal and rinse applicator thoroughly. 8. Refill bowl with water.

XXXX