



FORMULA 236

TERG-O-CIDE

**Disinfectant • Cleaner • Mildewstat
Fungicide • *Virucide • Deodorizer**

WHERE TO USE

Formula 236 Terg-O-Cide is recommended for use in hospitals, dental offices, health care facilities, nursing homes, daycare centers, correctional facilities, exercise facilities, locker rooms and schools.

WHAT TO USE ON

Formula 236 Terg-O-Cide is a multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 400ppm, in the presence of 5% soil load. It disinfects, cleans and deodorizes in one labor-saving step. It is effective on the following hard, non-porous surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, acrylic, etc.)



IS EFFECTIVE AGAINST

Formula 236 Terg-O-Cide is effective against: HIV-1 (AIDS virus), HBV and HCV on precleaned environmental surfaces/objects. It is also effective against Herpes Simplex 1 & Herpes Simplex 2, Athlete's Foot Fungus and Influenza A and Human Coronavirus. Canine parvovirus is no match for Formula 236 Terg-O-Cide at 8 oz. per gallon.

Care for Work Environments®

STATE FORMULA 236 TERG-O-CIDE

See hang tag for additional directions for use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARATION OF DISINFECTION/FUNGICIDAL/VIRUCIDAL USE SOLUTION: Apply 2 ounces per gallon (16 milliliters per liter) of water (1:64) to hard, inanimate, non-porous surfaces thoroughly wetting surfaces with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray. Treated surfaces must remain wet for 10 minutes. Rinse or allow to air dry. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use this product to clean or disinfect utensils, glassware, dishes or interior surfaces of appliances. Rinsing of floors is not necessary unless they are to be waxed or polished. Prepare a fresh solution at least daily or when use solution becomes diluted or soiled. Note: With spray applications cover or remove all food products.

For Disinfecting Hard, Non-Porous Surfaces Against Avian Influenza A: For use in homes, hospitals, hotels, motels and schools. The directions for hospital disinfection must be followed using a rate of 2 ounces per gallon of water. For use in animal housing facilities follow the animal premise directions.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

HOSPITAL/HEALTHCARE/INSTITUTIONAL

When used as directed at a 1:64 dilution (2 ounces per gallon of water), this product contains 660 ppm of active quaternary germicide, making it highly effective against a broad spectrum of pathogenic microorganisms.

Hepatitis Virus: At 2 ounces per gallon (660 ppm active quat) in the presence of 5% organic load this product was proven to be effective against (Duck) Hepatitis B Virus (HBV) and Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus) with a contact time of 10 minutes.

At 2 ounces per gallon of water in the presence of 400 ppm of hardness for a 10 minute contact time, this product was found to be effective against Hantavirus.

*KILLS HIV, HBV AND HCV ON PRECLEANED ENVIRONMENTAL

SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, (HIV-1) (associated with AIDS), (Duck) Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Clean up must always be done wearing protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** Leave surface wet for 10 minutes with a 2-ounce per gallon (16 milliliters per liter) use solution (or equivalent use dilution).

Mold and mildew control: To control the growth of mold and mildew and their odors on hard, non-porous surfaces, apply use solution to thoroughly wet all treated surfaces completely. Let surface air dry. Repeat application weekly or when growth or odor reappears. Use as directed at a 1:64 dilution (2 fluid ounces per gallon of water). Allow surface to remain wet for ten (10) minutes.

To control the growth of mold and mildew on large inflatable non-porous plastic and rubber structures (animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 2 ounces per gallon of water for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

For Disinfection and To Control the Growth of Mold and Mildew on Non-Porous Athletic Equipment:

For use on wrestling and gymnastic mats, athletic mats, exercise equipment, athletic training tables, physical therapy tables, athletic helmets, wrestling/boxing headgear, athletic shoe soles. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 2 ounces per gallon of water for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

Fungicidal Activity: At 2 ounces per gallon use-level this product is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) and Aspergillus niger on inanimate surfaces in locker rooms, dressing rooms, shower and bath areas and exercise facilities. Pre-clean all surfaces prior to using this product. Saturate surfaces with use solution for a period of 10 minutes. Allow solution to dry on surface and repeat treatment every seven days or when new growth appears.

TECHNICAL DATA

ACTIVE INGREDIENTS

Didecyl dimethyl ammonium chloride.....	2.54%
n-Alkyl (C1450%, C1240%, C1610%) dimethyl benzyl ammonium chloride.....	1.69%
OTHER INGREDIENT.....	95.77%
TOTAL.....	100.00%

EPA Reg. No. 10324-154-70799

Read complete label and MSDS before using product.

EPA EST. No. 70799-OH-1



State Industrial Products
3100 Hamilton Avenue
Cleveland, OH 44114
To Order Call: 1-866-747-2229
www.stateindustrial.com

Puerto Rico
Royal Industrial Park, Bldg "M"
Local #5, Carr 869, km 1.5 Palmas
Catano, PR. 00962
To Order Call: 1-800-782-2436

90R01
LIT800125

© Copyright 2009 — State Industrial Products Corp.